

Mercator Research Institute on
Global Commons and Climate Change



Spatial modeling of ecological-economic tradeoffs: renewable energy production & biodiversity conservation in the Alps

Sabine Fuss^{*,#} Florian Kraxner[#], Sylvain Leduc[#], Nicklas Forsell[#]

* Resources & International Trade, MCC, Berlin

Ecosystems Services & Management, IIASA, Laxenburg, Austria






Background

- Renewable energy (RE): EU climate change policy (2009/28/EC), nuclear phase-out
- Alpine Convention's Energy Protocol: Alpine region to make a long-term contribution to meeting Europe's energy needs (EC 2005, p. 37)
- Concern: ESS often compete with RE for productive sites \Rightarrow important tradeoffs to be analyzed to maintain ES functions and services under increasing RE demand and other pressures.
- Approach: use spatial optimization model to determine cost-optimal location of bioenergy plants under varying sustainability criteria (at different scales)
- Valuation concept: Analysis does not intend to assign monetary value to biodiversity and other ESS, but progressively excludes areas from optimization according to an ESS ranking \Rightarrow marginal costs of protecting those areas.
- Contributions:
 - Detailed spatial analysis of bioenergy solutions and tradeoffs in in multifunctional landscapes
 - Quantification and visualization of ecological-economic tradeoffs without making assumptions on weights and preferences and judgments about valuation
 - Aid decision-makers in forming strategies offering robustness across uncertainties




Marginal Protection Cost Concept

- ESS values difficult to quantify, need value judgements, surveys imperfect, no agreement on which ESS will be modeled, no consistent data sets, etc
- Suggested solution: Compute marginal protection costs much like the concept of marginal abatement costs:
 - circumvents the problems of assigning monetary value to ESS
 - enables us to say how much preservation will cost in terms of more expensive RE
 - leaving value judgements to preference of the user



Each additional ton of CO₂ abated will increase the cost of abating the next one, as low-cost opportunities are exploited and more expensive abatement options have to be employed.



Each additional area with important ESS excluded from RE deployment/protected will increase the cost of delivering another kWh of RE, as we have to resort to less productive areas.



Methodology

- Use the **BeWhere** Model* to optimize location and size of bioenergy units given the supply chain.
 - feedstock supply (incl. harvesting, transportation, etc.)
 - transportation routes/costs
 - trading possibilities
 - competition with other industries for feedstock
 - distribution networks
 - proximity to demand centers
- Mixed integer linear programming, economies of scale
- Geographically explicit on a 0.5 degree grid



*Leduc et al. (2012). CHP or biofuel production in Europe?
Energy Procedia, 20:40-49.



Technologies & Input Data

- Bioenergy technology: CHP, Gasification (43% heat / 42% power)
- Feedstocks: see map on next slide
- Bioenergy demand: based on national heat consumption (source: Werner 2006, Ecoheatcool WP1, WP4 / Berndes et al 2010)
- Transport networks, transportation cost (Source: National Imagery and Mapping Agency)
- Biodiversity hotspots / protected areas (Source: Econnect)





Biomass: current increment

Biomass: current stock





Source: G4M, forest inventory data







Bioenergy costs in the Alps with/out protection






Regional zoom: Vorarlberg's biomass






Vorarlberg's protected areas

- The sum of all protected/biodiverse areas still amounts to a substantial part of Vorarlberg's land area.
- But: reserves are often where biomass productivity is not the highest in the first place.





Marginal protection cost in Vorarlberg





Conclusions

- Alps:
 - If 20% of the biomass increment can be used for bioenergy, then approx. 20 PJ are used when all areas are protected, while up to 50 PJ are used if no areas are protected underlining the importance of analyzing the tradeoff between protection and bioenergy provision.
 - Low levels of bioenergy will be more expensive to generate if all biodiverse areas are protected, but the difference is marginal and will shrink as production expands and economies to scale are achieved.
- Vorarlberg:
 - Costs increase, as we move into areas where bioenergy generation is more expensive due to restrictions on location and scale of plants.
 - Marginal protection costs relatively low, especially for low production levels and not excluding regional parks.
 - Importance of zooming into regions, rather than drawing quick conclusions on aggregate estimates.



Contact & Acknowledgment

Sabine Fuss

Mercator Research Institute on
Global Commons and Climate
Change gGmbH

Torgauer Str. 12–15 | 10829
Berlin | Germany

mail fuss@mcc-berlin.net

web www.mcc-berlin.net

MCC was founded jointly by Stiftung Mercator
& Potsdam Institute for Climate Impact
Research



F. Kraxner & S. Leduc

ESM Program, International
Institute for Applied Systems
Analysis

Schlossplatz 1 | 2361
Laxenburg | Austria

Mail kraxner,leduc@iiasa.ac.at

web www.iiasa.ac.at

This research is part of an Alpine Space project
called recharge.green: <http://www.recharge-green.eu/>

